


What you should know about COVID-19 to protect yourself and others



Know about COVID-19

- Coronavirus (COVID-19) is an illness caused by a virus that can spread from person to person.
- The virus that causes COVID-19 is a new coronavirus that has spread throughout the world.
- COVID-19 symptoms can range from mild (or no symptoms) to severe illness.



Practice social distancing

- Buy groceries and medicine, go to the doctor, and complete banking activities online when possible.
- If you must go in person, stay at least 6 feet away from others and disinfect items you must touch.
- Get deliveries and takeout, and limit in-person contact as much as possible.



Know how COVID-19 is spread

- You can become infected by coming into close contact (about 6 feet or two arm lengths) with a person who has COVID-19. COVID-19 is primarily spread from person to person.
- You can become infected from respiratory droplets when an infected person coughs, sneezes, or talks.
- You may also be able to get it by touching a surface or object that has the virus on it, and then by touching your mouth, nose, or eyes.



Prevent the spread of COVID-19 if you are sick

- Stay home if you are sick, except to get medical care.
- Avoid public transportation, ride-sharing, or taxis.
- Separate yourself from other people and pets in your home.
- There is no specific treatment for COVID-19, but you can seek medical care to help relieve your symptoms.
- If you need medical attention, call ahead.



Protect yourself and others from COVID-19

- There is currently no vaccine to protect against COVID-19. The best way to protect yourself is to avoid being exposed to the virus that causes COVID-19.
- Stay home as much as possible and avoid close contact with others.
- Wear a cloth face covering that covers your nose and mouth in public settings.
- Clean and disinfect frequently touched surfaces.
- Wash your hands often with soap and water for at least 20 seconds, or use an alcohol-based hand sanitizer that contains at least 60% alcohol.



Know your risk for severe illness

- Everyone is at risk of getting COVID-19.
- Older adults and people of any age who have serious underlying medical conditions may be at higher risk for more severe illness.



Prevent the spread of COVID-19 if you are sick

Accessible version: <https://www.cdc.gov/coronavirus/2019-ncov/if-you-are-sick/steps-when-sick.html>

If you are sick with COVID-19 or think you might have COVID-19, follow the steps below to help protect other people in your home and community.

Stay home except to get medical care.

- **Stay home.** Most people with COVID-19 have mild illness and are able to recover at home without medical care. Do not leave your home, except to get medical care. Do not visit public areas.
- **Take care of yourself.** Get rest and stay hydrated.
- **Get medical care when needed.** Call your doctor before you go to their office for care. But, if you have trouble breathing or other concerning symptoms, call 911 for immediate help.
- **Avoid public transportation, ride-sharing, or taxis.**



Separate yourself from other people and pets in your home.

- **As much as possible, stay in a specific room** and away from other people and pets in your home. Also, you should use a separate bathroom, if available. If you need to be around other people or animals in or outside of the home, wear a cloth face covering.
 - See COVID-19 and Animals if you have questions about pets: <https://www.cdc.gov/coronavirus/2019-ncov/faq.html#COVID19animals>



Monitor your symptoms.

- **Common symptoms of COVID-19 include fever and cough.** Trouble breathing is a more serious symptom that means you should get medical attention.
- **Follow care instructions from your healthcare provider and local health department.** Your local health authorities will give instructions on checking your symptoms and reporting information.



If you develop **emergency warning signs** for COVID-19 get **medical attention immediately.**

Emergency warning signs include*:

- Trouble breathing
- Persistent pain or pressure in the chest
- New confusion or not able to be woken
- Bluish lips or face

*This list is not all inclusive. Please consult your medical provider for any other symptoms that are severe or concerning to you.

Call 911 if you have a medical emergency. If you have a medical emergency and need to call 911, notify the operator that you have or think you might have, COVID-19. If possible, put on a facemask before medical help arrives.

Call ahead before visiting your doctor.

- **Call ahead.** Many medical visits for routine care are being postponed or done by phone or telemedicine.
- **If you have a medical appointment that cannot be postponed, call your doctor's office.** This will help the office protect themselves and other patients.



If you are sick, wear a cloth covering over your nose and mouth.

- **You should wear a cloth face covering over your nose and mouth** if you must be around other people or animals, including pets (even at home).
- You don't need to wear the cloth face covering if you are alone. If you can't put on a cloth face covering (because of trouble breathing for example), cover your coughs and sneezes in some other way. Try to stay at least 6 feet away from other people. This will help protect the people around you.



Note: During the COVID-19 pandemic, medical grade facemasks are reserved for healthcare workers and some first responders. You may need to make a cloth face covering using a scarf or bandana.



[cdc.gov/coronavirus](https://www.cdc.gov/coronavirus)

Cover your coughs and sneezes.

- **Cover your mouth and nose** with a tissue when you cough or sneeze.
- **Throw used tissues** in a lined trash can.
- **Immediately wash your hands** with soap and water for at least 20 seconds. If soap and water are not available, clean your hands with an alcohol-based hand sanitizer that contains at least 60% alcohol.



Clean your hands often.

- **Wash your hands often** with soap and water for at least 20 seconds. This is especially important after blowing your nose, coughing, or sneezing; going to the bathroom; and before eating or preparing food.
- **Use hand sanitizer** if soap and water are not available. Use an alcohol-based hand sanitizer with at least 60% alcohol, covering all surfaces of your hands and rubbing them together until they feel dry.
- **Soap and water are the best option**, especially if your hands are visibly dirty.
- **Avoid touching** your eyes, nose, and mouth with unwashed hands.



Avoid sharing personal household items.

- **Do not share** dishes, drinking glasses, cups, eating utensils, towels, or bedding with other people in your home.
- **Wash these items thoroughly after using them** with soap and water or put them in the dishwasher.



Clean all “high-touch” surfaces everyday.

- **Clean and disinfect** high-touch surfaces in your “sick room” and bathroom. Let someone else clean and disinfect surfaces in common areas, but not your bedroom and bathroom.
- **If a caregiver or other person needs to clean and disinfect** a sick person’s bedroom or bathroom, they should do so on an as-needed basis. The caregiver/other person should wear a mask and wait as long as possible after the sick person has used the bathroom.



High-touch surfaces include phones, remote controls, counters, tabletops, doorknobs, bathroom fixtures, toilets, keyboards, tablets, and bedside tables.

- **Clean and disinfect areas that may have blood, stool, or body fluids on them.**


- **Use household cleaners and disinfectants.** Clean the area or item with soap and water or another detergent if it is dirty. Then use a household disinfectant.
 - Be sure to follow the instructions on the label to ensure safe and effective use of the product. Many products recommend keeping the surface wet for several minutes to ensure germs are killed. Many also recommend precautions such as wearing gloves and making sure you have good ventilation during use of the product.
 - Most EPA-registered household disinfectants should be effective.

How to discontinue home isolation

- People **with COVID-19 who have stayed home (home isolated)** can stop home isolation under the following conditions:
 - **If you will not have a test** to determine if you are still contagious, you can leave home after these three things have happened:
 - You have had no fever for at least 72 hours (that is three full days of no fever without the use of medicine that reduces fevers)AND
 - other symptoms have improved (for example, when your cough or shortness of breath has improved)AND
 - at least 7 days have passed since your symptoms first appeared.
 - **If you will be tested** to determine if you are still contagious, you can leave home after these three things have happened:
 - You no longer have a fever (without the use of medicine that reduces fevers)AND
 - other symptoms have improved (for example, when your cough or shortness of breath has improved)AND
 - you received two negative tests in a row, 24 hours apart. Your doctor will follow CDC guidelines.



In all cases, follow the guidance of your healthcare provider and local health department. The decision to stop home isolation should be made in consultation with your healthcare provider and state and local health departments. Local decisions depend on local circumstances.



Taking Care of Your Behavioral Health:

TIPS FOR SOCIAL DISTANCING, QUARANTINE, AND ISOLATION DURING AN INFECTIOUS DISEASE OUTBREAK

What Is Social Distancing?

Social distancing is a way to keep people from interacting closely or frequently enough to spread an infectious disease. Schools and other gathering places such as movie theaters may close, and sports events and religious services may be cancelled.

What Is Quarantine?

Quarantine separates and restricts the movement of people who have been exposed to a contagious disease to see if they become sick. It lasts long enough to ensure the person has not contracted an infectious disease.

What Is Isolation?

Isolation prevents the spread of an infectious disease by separating people who are sick from those who are not. It lasts as long as the disease is contagious.

Introduction

In the event of an infectious disease outbreak, local officials may require the public to take measures to limit and control the spread of the disease. This tip sheet provides information about **social distancing, quarantine, and isolation**. The government has the right to enforce federal and state laws related to public health if people

within the country get sick with highly contagious diseases that have the potential to develop into outbreaks or pandemics.

This tip sheet describes feelings and thoughts you may have during and after social distancing, quarantine, and isolation. It also suggests ways to care for your behavioral health during these experiences and provides resources for more help.

What To Expect: Typical Reactions

Everyone reacts differently to stressful situations such as an infectious disease outbreak that requires social distancing, quarantine, or isolation. People may feel:

- **Anxiety, worry, or fear** related to:
 - Your own health status
 - The health status of others whom you may have exposed to the disease
 - The resentment that your friends and family may feel if they need to go into quarantine as a result of contact with you
 - The experience of monitoring yourself, or being monitored by others for signs and symptoms of the disease
 - Time taken off from work and the potential loss of income and job security
 - The challenges of securing things you need, such as groceries and personal care items

- **Concern** about being able to effectively care for children or others in your care
 - **Uncertainty or frustration** about how long you will need to remain in this situation, and uncertainty about the future
 - **Loneliness** associated with feeling cut off from the world and from loved ones
 - **Anger** if you think you were exposed to the disease because of others' negligence
 - **Boredom and frustration** because you may not be able to work or engage in regular day-to-day activities
 - **Uncertainty or ambivalence** about the situation
 - **A desire to use alcohol or drugs** to cope
 - **Symptoms of depression**, such as feelings of hopelessness, changes in appetite, or sleeping too little or too much
 - **Symptoms of post-traumatic stress disorder (PTSD)**, such as intrusive distressing memories, flashbacks (reliving the event), nightmares, changes in thoughts and mood, and being easily startled
- Stay up to date on what is happening, while limiting your media exposure. Avoid watching or listening to news reports 24/7 since this tends to increase anxiety and worry. Remember that children are especially affected by what they hear and see on television.
 - Look to credible sources for information on the infectious disease outbreak (see page 3 for sources of reliable outbreak-related information).

BE YOUR OWN ADVOCATE

Speaking out about your needs is particularly important if you are in quarantine, since you may not be in a hospital or other facility where your basic needs are met. Ensure you have what you need to feel safe, secure, and comfortable.

- Work with local, state, or national health officials to find out how you can arrange for groceries and toiletries to be delivered to your home as needed.
- Inform health care providers or health authorities of any needed medications and work with them to ensure that you continue to receive those medications.

If you or a loved one experience any of these reactions for 2 to 4 weeks or more, contact your health care provider or one of the resources at the end of this tip sheet.

Ways To Support Yourself During Social Distancing, Quarantine, and Isolation

UNDERSTAND THE RISK

Consider the real risk of harm to yourself and others around you. The public perception of risk during a situation such as an infectious disease outbreak is often inaccurate. Media coverage may create the impression that people are in immediate danger when really the risk for infection may be very low. Take steps to get the facts:

EDUCATE YOURSELF

Health care providers and health authorities should provide information on the disease, its diagnosis, and treatment.

- Do not be afraid to ask questions—clear communication with a health care provider may help reduce any distress associated with social distancing, quarantine, or isolation.
- Ask for written information when available.
- Ask a family member or friend to obtain information in the event that you are unable to secure this information on your own.

WORK WITH YOUR EMPLOYER TO REDUCE FINANCIAL STRESS

If you're unable to work during this time, you may experience stress related to your job status or financial situation.

- Provide your employer with a clear explanation of why you are away from work.
- Contact the U.S. Department of Labor toll-free at 1-866-4USWAGE (1-866-487-9243) about the Family and Medical Leave Act (FMLA), which allows U.S. employees up to 12 weeks of unpaid leave for serious medical conditions, or to care for a family member with a serious medical condition.
- Contact your utility providers, cable and Internet provider, and other companies from whom you get monthly bills to explain your situation and request alternative bill payment arrangements as needed.
- If approved by health authorities and your health care providers, arrange for your friends and loved ones to bring you newspapers, movies, and books.
- Sign up for emergency alerts via text or email to ensure you get updates as soon as they are available.
- Call SAMHSA's free 24-hour Disaster Distress Helpline at 1-800-985-5990, if you feel lonely or need support.
- Use the Internet, radio, and television to keep up with local, national, and world events.
- If you need to connect with someone because of an ongoing alcohol or drug problem, consider calling your local Alcoholics Anonymous or Narcotics Anonymous offices.

Sources for Reliable Outbreak-Related Information

Centers for Disease Control and Prevention

1600 Clifton Road
Atlanta, GA 30329-4027
1-800-CDC-INFO (1-800-232-4636)
<http://www.cdc.gov>

World Health Organization

Regional Office for the Americas of the World Health Organization
525 23rd Street, NW
Washington, DC 20037
202-974-3000
<http://www.who.int/en>

CONNECT WITH OTHERS

Reaching out to people you trust is one of the best ways to reduce anxiety, depression, loneliness, and boredom during social distancing, quarantine, and isolation. You can:

- Use the telephone, email, text messaging, and social media to connect with friends, family, and others.
- Talk “face to face” with friends and loved ones using Skype or FaceTime.

TALK TO YOUR DOCTOR

If you are in a medical facility, you may have access to health care providers who can answer your questions. However, if you are quarantined at home, and you're worried about physical symptoms you or your loved ones may be experiencing, call your doctor or other health care provider:

- Ask your provider whether it would be possible to schedule remote appointments via Skype or FaceTime for mental health, substance use, or physical health needs.
- In the event that your doctor is unavailable and you are feeling stressed or are in crisis, call the hotline numbers listed at the end of this tip sheet for support.

USE PRACTICAL WAYS TO COPE AND RELAX

- Relax your body often by doing things that work for you—take deep breaths, stretch, meditate or pray, or engage in activities you enjoy.
- Pace yourself between stressful activities, and do something fun after a hard task.

- Talk about your experiences and feelings to loved ones and friends, if you find it helpful.
- Maintain a sense of hope and positive thinking; consider keeping a journal where you write down things you are grateful for or that are going well.

After Social Distancing, Quarantine, or Isolation

You may experience mixed emotions, including a sense of relief. If you were isolated because you had the illness, you may feel sadness or anger because friends and loved ones may have unfounded fears of contracting the disease from contact with you, even though you have been determined not to be contagious.

The best way to end this common fear is to learn about the disease and the actual risk to others. Sharing this information will often calm fears in others and allow you to reconnect with them.

If you or your loved ones experience symptoms of extreme stress—such as trouble sleeping, problems with eating too much or too little, inability to carry out routine daily activities, or using drugs or alcohol to cope—speak to a health care provider or call one of the hotlines listed to the right for a referral.

If you are feeling overwhelmed with emotions such as sadness, depression, anxiety, or feel like you want to harm yourself or someone else, call 911 or the National Suicide Prevention Lifeline at 1-800-273-TALK (1-800-273-8255).

Helpful Resources

Hotlines

For Cattaraugus County:

OGH crisis hotline:

1-800-339-5209

County crisis hotline:

716-373-8040

SAMHSA's Disaster Distress Helpline

Toll-Free: 1-800-985-5990 (English and español)

SMS: Text TalkWithUs to 66746

SMS (español): "Hablamos" al 66746

TTY: 1-800-846-8517

Website (English): <http://www.disasterdistress.samhsa.gov>

Website (español): <http://www.disasterdistress.samhsa.gov/espanol.aspx>

SAMHSA's National Helpline

Toll-Free: 1-800-662-HELP (24/7/365 Treatment Referral Information Service in English and español)

Website: <http://www.samhsa.gov/find-help/national-helpline>

National Suicide Prevention Lifeline

Toll-Free (English): 1-800-273-TALK (8255)

Toll-Free (español): 1-888-628-9454

TTY: 1-800-799-4TTY (4889)

Website (English): <http://www.suicidepreventionlifeline.org>

Website (español): <http://www.suicidepreventionlifeline.org/gethelp/spanish.aspx>

Treatment Locator

Behavioral Health Treatment Services Locator

Website: <http://findtreatment.samhsa.gov/locator/home>

**Note: Inclusion or mention of a resource in this fact sheet does not imply endorsement by the Center for Mental Health Services, the Substance Abuse and Mental Health Services Administration, or the U.S. Department of Health and Human Services.*



HHS Publication No. SMA-14-4894
(2014)